arXiv:1701.08929v1 [math.AP] 31 Jan 2017

FACTORIZATIONS AND HARDY–RELLICH-TYPE INEQUALITIES

FRITZ GESZTESY AND LANCE LITTLEJOHN

Dedicated with great pleasure to Helge Holden on the occasion of his 60th birthday.

Abstract. The principal aim of this note is to illustrate how factorizations of singular, even-order partial differential operators yield an elementary approach to classical inequalities of Hardy–Rellich-type. More precisely, introducing the two-parameter n-dimensional homogeneous scalar differential expressions $T_{\alpha,\beta} := -\Delta + \alpha|x|^{-2} x\cdot\nabla + \beta|x|^{-2}$, $\alpha,\beta\in\mathbb{R}$, $x\in\mathbb{R}^n\backslash\{0\}$, $n\in\mathbb{N}$, $n\geq 2$, and its formal adjoint, denoted by $T^+_{\alpha,\beta}$, we show that nonnegativity of $T^+_{\alpha,\beta}T_{\alpha,\beta}$ on $C_0^\infty(\mathbb{R}^n\backslash\{0\})$ implies the fundamental inequality,

$$\begin{aligned}\int_{\mathbb{R}^n}[(\Delta f)(x)]^2\,d^nx &\geq [(n-4)\alpha-2\beta]\int_{\mathbb{R}^n}|x|^{-2}|(\nabla f)(x)|^2\,d^nx\\ &\quad-\alpha(\alpha-4)\int_{\mathbb{R}^n}|x|^{-4}|x\cdot(\nabla f)(x)|^2\,d^nx\\ &\quad+\beta[(n-4)(\alpha-2)-\beta]\int_{\mathbb{R}^n}|x|^{-4}|f(x)|^2\,d^nx,\quad f\in C_0^\infty(\mathbb{R}^n\backslash\{0\}).\end{aligned}\tag{0.1}$$

A particular choice of values for α and β in (0.1) yields known Hardy–Rellich-type inequalities, including the classical Rellich inequality and an inequality due to Schmincke. By locality, these inequalities extend to the situation where $\mathbb{R}^n$ is replaced by an arbitrary open set $\Omega\subseteq\mathbb{R}^n$ for functions $f\in C_0^\infty(\Omega\backslash\{0\})$.

Perhaps more importantly, we will indicate that our method, in addition to being elementary, is quite flexible when it comes to a variety of generalized situations involving the inclusion of remainder terms and higher-order situations.

1. Introduction

We dedicate this note with great pleasure to Helge Holden, whose wide range of contributions to a remarkable variety of areas in mathematical physics, stochastics, partial differential equations, and integrable systems, whose exemplary involvement with students, and whose tireless efforts on behalf of the mathematical community, deserve our utmost respect and admiration. Happy Birthday, Helge, we hope our modest contribution to Hardy–Rellich-type inequalities will give some joy.

The celebrated (multi-dimensional) Hardy inequality,

$$\int_{\mathbb{R}^n}|(\nabla f)(x)|^2\,d^nx\geq[(n-2)/2]^2\int_{\mathbb{R}^n}|x|^{-2}|f(x)|^2\,d^nx,\quad f\in C_0^\infty(\mathbb{R}^n\backslash\{0\}),\ n\in\mathbb{N},\ n\geq 3,\tag{1.1}$$

and Rellich's inequality,

$$\int_{\mathbb{R}^n}|(\Delta f)(x)|^2\,d^nx\geq[n(n-4)/4]^2\int_{\mathbb{R}^n}|x|^{-4}|f(x)|^2\,d^nx,\quad f\in C_0^\infty(\mathbb{R}^n\backslash\{0\}),\ n\in\mathbb{N},\ n\geq 5,\tag{1.2}$$

Date: February 1, 2017.

2010 *Mathematics Subject Classification.* Primary: 35A23, 35J30; Secondary: 47A63, 47F05.

Key words and phrases. Hardy inequality, Rellich-type inequality, factorizations of differential operators.

the first two inequalities in an infinite sequence of higher-order Hardy-type inequalities, received enormous attention in the literature due to their ubiquity in self-adjointness and spectral theory problems associated with second and fourth-order differential operators with strongly singular coefficients, respectively (see, e.g., [2], [3], [6], [10], [15, Sect. 1.5], [16, Ch. 5], [28], [31], [32], [35], [38]–[42], [54, Ch. II], [57]). We refer to Remark 2.11 for a selection of Rellich inequality references and some pertinent monographs on Hardy's inequality.

As one of our principal results we will derive the following two-parameter family of inequalities (a special case of inequality (1.5) below): If either $\alpha \leq 0$ or $\alpha \geq 4$, and $\beta \in \mathbb{R}$, then,

$$\begin{aligned}\int_{\mathbb{R}^n} |(\Delta f)(x)|^2 \, d^n x \geq [\alpha(n-\alpha) - 2\beta] \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2 \, d^n x \\ + \beta[(n-4)(\alpha-2) - \beta] \int_{\mathbb{R}^n} |x|^{-4} |f(x)|^2 \, d^n x, \\ f \in C_0^\infty(\mathbb{R}^n \backslash \{0\}), \, n \in \mathbb{N}, \, n \geq 2.\end{aligned} \tag{1.3}$$

As will be shown, (1.3) contains Rellich's inequality (1.2), and Schmincke's one-parameter family of inequalities,

$$\begin{aligned}\int_{\mathbb{R}^n} |(\Delta f)(x)|^2 \, d^n x \geq -s \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2 \, d^n x \\ + [(n-4)/4]^2 \big(4s + n^2\big) \int_{\mathbb{R}^n} |x|^{-4} |f(x)|^2 \, d^n x, \\ s \in \big[-2^{-1} n(n-4), \infty \big),\end{aligned} \tag{1.4}$$

as special cases. By locality, the inequalities (1.1)–(1.4) naturally extend to the case where $\mathbb{R}^n$ is replaced by an arbitrary open set $\Omega \subset \mathbb{R}^n$ for functions $f \in C_0^\infty(\Omega \backslash \{0\})$ (without changing the constants in these inequalities).

Our approach is based on factorizing even-order differential equations. More precisely, focusing on the 4th-order case for simplicity, we introduce the two-parameter n-dimensional homogeneous scalar differential expressions $T_{\alpha,\beta} := -\Delta + \alpha |x|^{-2} x \cdot \nabla + \beta |x|^{-2}$, $\alpha, \beta \in \mathbb{R}$, $x \in \mathbb{R}^n \backslash \{0\}$, $n \in \mathbb{N}$, $n \geq 2$, and its formal adjoint, denoted by $T_{\alpha,\beta}^+$. Nonnegativity of $T_{\alpha,\beta}^+ T_{\alpha,\beta}$ on $C_0^\infty(\mathbb{R}^n \backslash \{0\})$ then implies the fundamental inequality,

$$\begin{aligned}\int_{\mathbb{R}^n} [(\Delta f)(x)]^2 \, d^n x \geq [(n-4)\alpha - 2\beta] \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2 \, d^n x \\ - \alpha(\alpha-4) \int_{\mathbb{R}^n} |x|^{-4} |x \cdot (\nabla f)(x)|^2 \, d^n x \\ + \beta[(n-4)(\alpha-2) - \beta] \int_{\mathbb{R}^n} |x|^{-4} |f(x)|^2 \, d^n x, \quad f \in C_0^\infty(\mathbb{R}^n \backslash \{0\}),\end{aligned} \tag{1.5}$$

which in turn contains inequality (1.3) as a special case.

We conclude our note with a series of remarks putting our approach into proper context by indicating that our method is elementary and very flexible in handling a variety of generalized situations involving the inclusion of remainder terms and higher even-order differential operators.

2. Factorizations and Hardy–Rellich-type Inequalities

The principal inequality to be proven in this note is of the following form:

Theorem 2.1. *Let $\alpha, \beta \in \mathbb{R}$, and $f \in C_0^\infty(\mathbb{R}^n\backslash\{0\})$, $n \in \mathbb{N}$, $n \geq 2$. Then,*

$$\begin{aligned}\int_{\mathbb{R}^n} [(\Delta f)(x)]^2 \, d^n x &\geq [(n-4)\alpha - 2\beta] \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2 \, d^n x \\ &\quad - \alpha(\alpha-4) \int_{\mathbb{R}^n} |x|^{-4} |x \cdot (\nabla f)(x)|^2 \, d^n x \\ &\quad + \beta[(n-4)(\alpha-2) - \beta] \int_{\mathbb{R}^n} |x|^{-4} |f(x)|^2 \, d^n x. \end{aligned} \tag{2.1}$$

In addition, if either $\alpha \leq 0$ or $\alpha \geq 4$, then,

$$\begin{aligned}\int_{\mathbb{R}^n} |(\Delta f)(x)|^2 \, d^n x &\geq [\alpha(n-\alpha) - 2\beta] \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2 \, d^n x \\ &\quad + \beta[(n-4)(\alpha-2) - \beta] \int_{\mathbb{R}^n} |x|^{-4} |f(x)|^2 \, d^n x. \end{aligned} \tag{2.2}$$

Proof. Given $\alpha, \beta \in \mathbb{R}$ and $n \in \mathbb{N}$, $n \geq 2$, we introduce the two-parameter n-dimensional homogeneous scalar differential expressions

$$T_{\alpha,\beta} := -\Delta + \alpha |x|^{-2} x \cdot \nabla + \beta |x|^{-2}, \quad x \in \mathbb{R}^n\backslash\{0\}, \tag{2.3}$$

and its formal adjoint, denoted by $T_{\alpha,\beta}^+$,

$$T_{\alpha,\beta}^+ := -\Delta - \alpha |x|^{-2} x \cdot \nabla + [\beta - \alpha(n-2)] |x|^{-2}, \quad x \in \mathbb{R}^n\backslash\{0\}. \tag{2.4}$$

Assuming $f \in C_0^\infty(\mathbb{R}^n\backslash\{0\})$ throughout this proof, employing elementary multi-variable differential calculus, we proceed to the computation of $T_{\alpha,\beta}^+ T_{\alpha,\beta}$ (which, while entirely straightforward, may well produce some tears in the process),

$$\begin{aligned}(T_{\alpha,\beta}^+ T_{\alpha,\beta} f)(x) &= (\Delta^2 f)(x) + [(n-4)\alpha - 2\beta] |x|^{-2} (\Delta f)(x) \\ &\quad + \alpha(4-\alpha) |x|^{-4} \sum_{j,k=1}^n x_j x_k f_{x_j,x_k}(x) \\ &\quad + \big[-(n-3)\alpha^2 + 2(n-2)\alpha + 4\beta \big] |x|^{-4} x \cdot (\nabla f)(x) \\ &\quad + \big[\beta^2 + 2(n-4)\beta - (n-4)\alpha\beta\big] |x|^{-4} f(x). \end{aligned} \tag{2.5}$$

Thus, choosing $f \in C_0^\infty(\mathbb{R}^n\backslash\{0\})$ real-valued from this point on and integrating by parts (observing the support properties of f, which results in vanishing surface terms) implies

$$\begin{aligned} 0 &\leq \int_{\mathbb{R}^n} [(T_{\alpha,\beta} f)(x)]^2 \, d^n x = \int_{\mathbb{R}^n} f(x) (T_{\alpha,\beta}^+ T_{\alpha,\beta} f)(x) \, d^n x \\ &= \int_{\mathbb{R}^n} [(\Delta f)(x)]^2 \, d^n x + [(n-4)\alpha - 2\beta] \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |x|^{-2} f(x) (\Delta f)(x) \, d^n x \\ &\quad + \alpha(\alpha-4) \sum_{j,k=1}^n \int_{\mathbb{R}^n} |x|^{-4} f(x) x_j x_k f_{x_j,x_k}(x) \, d^n x \\ &\quad + \big[-(n-3)\alpha^2 + 2(n-2)\alpha + 4\beta \big] \int_{\mathbb{R}^n} |x|^{-4} f(x) [x \cdot (\nabla f)(x)] \, d^n x \\ &\quad + \big[\beta^2 + 2(n-4)\beta - (n-4)\alpha\beta\big] \int_{\mathbb{R}^n} |x|^{-4} f(x)^2 \, d^n x. \end{aligned} \tag{2.6}$$

To simplify and exploit expression (2.6), we make two observations. First, a standard integration by parts (again observing the support properties of f) yields

$$\begin{aligned}\int_{\mathbb{R}^n} |x|^{-2} f(x)(\Delta f)(x)\, d^n x = 2\int_{\mathbb{R}^n} |x|^{-4} f(x)(x\cdot(\nabla f)(x)\, d^n x \\ - \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2\, d^n x.\end{aligned} \tag{2.7}$$

Similarly, one confirms that

$$\begin{aligned}\sum_{j,k=1}^n \int_{\mathbb{R}^n} x_j x_k f(x) f_{x_j,x_k}(x) = -(n-3)\int_{\mathbb{R}^n} |x|^{-4} f(x)[x\cdot(\nabla f)(x)]\, d^n x \\ - \int_{\mathbb{R}^n} |x|^{-4}[x\cdot(\nabla f)(x)]^2\, d^n x.\end{aligned} \tag{2.8}$$

Combining (2.6)–(2.8) then yields (2.1).

Since by Cauchy's inequality,

$$-\int_{\mathbb{R}^n} |x|^{-4}[x\cdot(\nabla f)(x)]^2\, d^n x \geq -\int_{\mathbb{R}^n} |x|^{-2}|(\nabla f)(x)|^2\, d^n x, \tag{2.9}$$

one concludes that as long as $\alpha(\alpha-4)\geq 0$, that is, as long as either $\alpha\leq 0$ or $\alpha\geq 4$, one can further estimate (2.1) from below and thus arrive at inequality (2.2). □

As a special case of (2.2) one obtains Rellich's classical inequality in its original form as follows:

Corollary 2.2. *Let $n\in\mathbb{N}$, $n\geq 5$, and $f\in C_0^\infty(\mathbb{R}^n\backslash\{0\})$. Then,*

$$\int_{\mathbb{R}^n} |(\Delta f)(x)|^2\, d^n x \geq [n(n-4)/4]^2 \int_{\mathbb{R}^n} |x|^{-4}|f(x)|^2\, d^n x. \tag{2.10}$$

Proof. Choosing $\beta=\alpha(n-\alpha)/2$ in (2.2) results in

$$\int_{\mathbb{R}^n} |(\Delta f)(x)|^2\, d^n x \geq G_n(\alpha)\int_{\mathbb{R}^n} |x|^{-4}|f(x)|^2\, d^n x, \tag{2.11}$$

with

$$G_n(\alpha)=\alpha(n-\alpha)\{(n-4)(\alpha-2)-[\alpha(n-\alpha)/2]\}/2. \tag{2.12}$$

Maximizing $G_n(\alpha)$ with respect to α (it is advantageous to introduce the new variable $a=\alpha-2$) yields maxima at

$$\alpha_\pm = 2\pm\big[\big(n^2/2\big)-2n+4\big]^{1/2}, \tag{2.13}$$

and taking the constraints $\alpha\leq 0$ or $\alpha\geq 4$ into account results in $n\geq 5$. The fact

$$G_n(\alpha_\pm)=[n(n-4)/4]^2, \tag{2.14}$$

then yields the Rellich inequality (2.10). □

Inequality (2.1) also implies the following result:

Corollary 2.3. *Let $n\in\mathbb{N}$ and $f\in C_0^\infty(\mathbb{R}^n\backslash\{0\})$. Then,*

$$\int_{\mathbb{R}^n} |(\Delta f)(x)|^2\, d^n x \geq \big(n^2/4\big)\int_{\mathbb{R}^n} |x|^{-2}|(\nabla f)(x)|^2\, d^n x, \quad n\geq 8, \tag{2.15}$$

and

$$\int_{\mathbb{R}^n} |(\Delta f)(x)|^2\, d^n x \geq 4(n-4)\int_{\mathbb{R}^n} |x|^{-2}|(\nabla f)(x)|^2\, d^n x, \quad 5\leq n\leq 7. \tag{2.16}$$

In addition,

$$\int_{\mathbb{R}^n} |(\Delta f)(x)|^2 \, d^n x \geq \big(n^2/4\big) \int_{\mathbb{R}^n} |x|^{-4} |x \cdot (\nabla f)(x)|^2 \, d^n x, \quad n \geq 2. \tag{2.17}$$

Proof. Again, we chose $f \in C_0^\infty(\mathbb{R}^n \backslash \{0\})$ real-valued for simplicity throughout this proof. The choice $\beta = (n-4)(\alpha-2)$ in (2.1) then results in

$$\begin{aligned} \int_{\mathbb{R}^n} |(\Delta f)(x)|^2 \, d^n x \geq (n-4)(4-\alpha) \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2 \, d^n x \\ - \alpha(\alpha-4) \int_{\mathbb{R}^n} |x|^{-4} |x \cdot (\nabla f)(x)|^2 \, d^n x. \end{aligned} \tag{2.18}$$

If in addition $\alpha < 0$, then applying Cauchy's inequality to the 2nd term on the right-hand side of (2.18) yields

$$\int_{\mathbb{R}^n} |(\Delta f)(x)|^2 \, d^n x \geq H_n(\alpha) \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2 \, d^n x, \tag{2.19}$$

where $H_n(\alpha) = (n-4+\alpha)(4-\alpha)$. Maximizing H_n with respect to α subject to the constraint $\alpha < 0$ yields a maximum at $\alpha_1 = (8-n)/2$, with $H_n((8-n)/2) = n^2/4$, implying inequality (2.15) for $n \geq 9$. On the other hand, choosing $\alpha = 0$ in (2.18) yields

$$\int_{\mathbb{R}^n} |(\Delta f)(x)|^2 \, d^n x \geq 4(n-4) \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2 \, d^n x. \tag{2.20}$$

Since $4(n-4) = n^2/4$ for $n = 8$, this proves (2.15). Actually, one can arrive at (2.15) much quicker, but since we will subsequently use (2.18), we kept the above argument in this proof: Indeed, choosing $\beta = 0$ in (2.2) yields

$$\int_{\mathbb{R}^n} |(\Delta f)(x)|^2 \, d^n x \geq \alpha(n-\alpha) \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2 \, d^n x. \tag{2.21}$$

Maximizing $F_n(\alpha) = \alpha(n-\alpha)$ with respect to α yields a maximum at $\alpha_1 = n/2$, and subjecting it to the constraint $\alpha \geq 4$ proves (2.15).

Choosing $\alpha = 4$, $\beta = 0$ in (2.1) yields (2.16).

For $n \geq 2$ and $(4-n) < \alpha < 4$, applying Cauchy's inequality to the 1st term on the right-hand side of (2.18) now yields

$$\int_{\mathbb{R}^n} |(\Delta f)(x)|^2 \, d^n x \geq K_n(\alpha) \int_{\mathbb{R}^n} |x|^{-4} |x \cdot (\nabla f)(x)|^2 \, d^n x, \tag{2.22}$$

where $K_n(\alpha) = -(\alpha+n-4)(\alpha-4)$. Maximizing K_n subject to the constraint $(4-n) < \alpha < 4$ yields a maximum at $\alpha_1 = (8-n)/2$, with $K_n((8-n)/2) = n^2/4$, implying (2.17). □

We conclude with a series of remarks that put our approach into proper context and point out natural continuations into various other directions.

Remark 2.4. (*i*) The constant in inequality (2.10) is known to be optimal, see, for instance, [6, p. 222], [17], [47], [50], [58], [61].
(*ii*) A sequence of extensions of (2.15), valid for $n \geq 5$, and for bounded domains containing 0, was derived by Tertikas and Zographopoulos [58, Theorem 1.7]. Moreover, an extension of inequality (2.15) valid for $n = 4$ and for bounded open domains containing 0 was proved by [1, Theorem 2.1 (b)]. An alternative inequality whose special cases also imply Rellich's inequality (2.10) and inequality (2.15) appeared in [14]. Thus, while the constant $n^2/4$ in (2.15) is known to be optimal (cf. [58] for $n \geq 5$), the constant $4(n-4)$ in (2.16) is not, the sharp constant being known to be $n^2/4$ (also for $n = 4$, cf. [1]). ⋄

Next, we comment on a special case of inequality (2.2) originally due to Schmincke [56]:

Remark 2.5. The choice $\beta = 2^{-1}(n-4)[\alpha - 2 - 4^{-1}(n-4)]$, and the introduction of the new variable

$$s = s(\alpha) = \alpha^2 - 4\alpha - 2^{-1}n(n-4), \tag{2.23}$$

renders the two-parameter inequality (2.2) into Schmincke's one-parameter inequality

$$\begin{aligned}\int_{\mathbb{R}^n} |(\Delta f)(x)|^2\, d^n x \geq -s \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2\, d^n x \\ + [(n-4)/4]^2 \big(4s + n^2\big) \int_{\mathbb{R}^n} |x|^{-4} |f(x)|^2\, d^n x, \\ s \in \big[-2^{-1}n(n-4), \infty\big).\end{aligned} \tag{2.24}$$

Here the requirements $\alpha \leq 0$, equivalently, $\alpha \geq 4$, both yield the range requirement for s in the form $s \in \big[-2^{-1}n(n-4), \infty\big)$. Inequality (2.24) is precisely the content of Lemma 2 in Schmincke [56], in particular, (2.2) thus recovers Schmincke's result. Moreover, assuming $n \geq 5$ (the case $n = 4$ being trivial) permits the value $s = 0$ and hence implies Rellich's inequality (2.10). If $n \geq 8$, the value $s = -n^2/4$ is permitted, yielding inequality (2.15). Finally, for $5 \leq n \leq 7$, $s \in \big[-2^{-1}n(n-4), \infty\big)$ and $4s + n^2 \geq 0$ permit one to choose $s = -n(n-4)/2$ and hence to conclude

$$\begin{aligned}\int_{\mathbb{R}^n} |(\Delta f)(x)|^2\, d^n x \geq 2^{-1}n(n-4) \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2\, d^n x \\ + [(n-4)/4]^2 \big(4s + n^2\big) \int_{\mathbb{R}^n} |x|^{-4} |f(x)|^2\, d^n x (2.21)\end{aligned} \tag{2.25}$$

$$\geq 2^{-1}n(n-4) \int_{\mathbb{R}^n} |x|^{-2} |(\nabla f)(x)|^2\, d^n x, \quad 5 \leq n \leq 7, \tag{2.26}$$

but inequality (2.16) is strictly superior to (2.26). Hence the two-parameter version (2.2) yields the better result (2.16), even though, the latter is not optimal either as mentioned in the previous Remark 2.4 (ii). $\diamond$

Remark 2.6. Since all differential expressions employed are local, and only integration by parts was involved in deriving (2.6) (cf., e.g., [39, Remark 4] in this context), the estimates (2.1), (2.2), (2.10), (2.15)–(2.17), (2.24), all extend to the case where $\mathbb{R}^n$ is replaced by an arbitrary open set $\Omega \subset \mathbb{R}^n$ for functions $f \in C_0^\infty(\Omega\backslash\{0\})$ (without changing the constants in these inequalities). $\diamond$

Remark 2.7. Since $C_0^\infty(\mathbb{R}^n\backslash\{0\})$ is dense in $H^2(\mathbb{R}^n)$ if and only if $n \geq 4$, equivalently, $-\Delta\big|_{C_0^\infty(\mathbb{R}^n\backslash\{0\})}$ is essentially self-adjoint in $L^2(\mathbb{R}^n)$ if and only if $n \geq 4$ (see, e.g., [21, p. 412–413], see also [41], [56]), Rellich's inequality (2.10) extends from $C_0^\infty(\mathbb{R}^n\backslash\{0\})$ to $H^2(\mathbb{R}^n)$ for $n \geq 5$, and inequalities (2.15), and (2.17) extend from $C_0^\infty(\mathbb{R}^n\backslash\{0\})$ to $H^2(\mathbb{R}^n)$ for $n \geq 4$. $\diamond$

Remark 2.8. This factorization approach was originally employed in the context of the classical Hardy inequality in [31] (and some of its logarithmic refinements in [28]). Without repeating the analogous steps in detail we just mention that given $n \in \mathbb{N}$, $n \geq 3$, $\alpha \in \mathbb{R}$, one introduces the one-parameter family of homogeneous vector-valued differential expressions

$$T_\alpha := \nabla + \alpha |x|^{-2} x, \quad x \in \mathbb{R}^n\backslash\{0\}, \tag{2.27}$$

with formal adjoint, denoted by T_α^+,

$$T_\alpha^+ = -\mathrm{div}(\,\cdot\,) + \alpha |x|^{-2} x\,\cdot, \quad x \in \mathbb{R}^n\backslash\{0\}, \tag{2.28}$$

such that (e.g., on $C_0^\infty(\mathbb{R}^n\backslash\{0\})$-functions),

$$T_\alpha^+ T_\alpha = -\Delta + \alpha(\alpha + 2 - n)|x|^{-2}. \tag{2.29}$$

Thus, for $f \in C_0^\infty(\mathbb{R}^n\backslash\{0\})$,

$$\begin{aligned} 0 &\leq \int_{\mathbb{R}^n} |T_\alpha f)(x)|^2 \, d^n x = \int_{\mathbb{R}^n} \overline{f(x)} (T_\alpha^+ T_\alpha f)(x) \, d^n x \\ &= \int_{\mathbb{R}^n} |(\nabla f)(x)|^2 \, d^n x + \alpha(\alpha + 2 - n) \int_{\mathbb{R}^n} |x|^{-2} |f(x)|^2 \, d^n x, \end{aligned} \tag{2.30}$$

and hence,

$$\int_{\mathbb{R}^n} |(\nabla f)(x)|^2 \, d^n x \geq \alpha[(n-2) - \alpha] \int_{\mathbb{R}^n} |x|^{-2} |f(x)|^2 \, d^n x. \tag{2.31}$$

Maximizing $\alpha[(n-2)-\alpha]$ with respect to α yields the classical Hardy inequality,

$$\int_{\mathbb{R}^n} |(\nabla f)(x)|^2 \, d^n x \geq [(n-2)/2]^2 \int_{\mathbb{R}^n} |x|^{-2} |f(x)|^2 \, d^n x, \quad f \in C_0^\infty(\mathbb{R}^n\backslash\{0\}). \tag{2.32}$$

Again, it is well-known that the constant in (2.32) is optimal (cf., e.g., [61]). ⋄

Actually, our factorization approach also yields a known improvement of Hardy's inequality (see, e.g., [6, Theorem 1.2.5], specializing it to $p = 2$, $\varepsilon = 0$). Next, we briefly sketch the corresponding argument.

Remark 2.9. Given $n \in \mathbb{N}$, $n \geq 3$, $\alpha \in \mathbb{R}$, one introduces the following modified one-parameter family of homogeneous vector-valued differential expressions

$$\widetilde{T}_\alpha := \big(|x|^{-1}x\big) \cdot \nabla + \alpha |x|^{-1}, \quad x \in \mathbb{R}^n\backslash\{0\}, \tag{2.33}$$

with formal adjoint, denoted by $\big(\widetilde{T}_\alpha\big)^+$,

$$\big(\widetilde{T}_\alpha\big)^+ = -\big(|x|^{-1}x\big) \cdot \nabla + (\alpha - n + 1)|x|^{-1}, \quad x \in \mathbb{R}^n\backslash\{0\}. \tag{2.34}$$

Exploiting the identities (for $f \in C_0^\infty(\mathbb{R}^n\backslash\{0\})$, for simplicity),

$$\big[|x|^{-1}x \cdot \nabla\big]\big[|x|^{-1}x \cdot (\nabla f)(x)\big] = |x|^{-2} \sum_{j,k=1}^n x_j x_k f_{x_j,x_k}(x), \quad x \in \mathbb{R}^n\backslash\{0\}, \tag{2.35}$$

$$x \cdot \nabla\big(|x|^{-1} f(x)\big) = |x|^{-1}[x \cdot (\nabla f)(x)] - |x|^{-1} f(x), \quad x \in \mathbb{R}^n\backslash\{0\}, \tag{2.36}$$

one computes (e.g., on $C_0^\infty(\mathbb{R}^n\backslash\{0\})$-functions),

$$\big(\widetilde{T}_\alpha\big)^+ \widetilde{T}_\alpha = -|x|^{-2} \sum_{j,k=1}^n x_j x_k \partial_{x_j} \partial_{x_k} - (n-1)|x|^{-2}[x \cdot (\nabla f)(x)] + \alpha(\alpha + 2 - n)|x|^{-2}, \quad x \in \mathbb{R}^n\backslash\{0\}. \tag{2.37}$$

Thus, appropriate integration by parts yield

$$\begin{aligned} 0 &\leq \int_{\mathbb{R}^n} \big|\big(\widetilde{T}_\alpha f\big)(x)\big|^2 \, d^n x = \int_{\mathbb{R}^n} \overline{f(x)} \big(\big(\widetilde{T}_\alpha\big)^+ \widetilde{T}_\alpha f\big)(x) \, d^n x \\ &= -\int_{\mathbb{R}^n} |x|^{-2} \bigg\{ \sum_{j,k=1}^n x_j x_k \overline{f(x)} f_{x_j,x_k}(x) + (n-1)\overline{f(x)}[x \cdot (\nabla f)(x)] \\ &\qquad - \alpha(\alpha - n + 2)|f(x)|^2 \bigg\} d^n x, \\ &= \int_{\mathbb{R}^n} |x|^{-2} \Big\{ |[x \cdot (\nabla f)(x)]|^2 + \alpha(\alpha - n + 2)|f(x)|^2 \Big\}, \quad f \in C_0^\infty(\mathbb{R}^n\backslash\{0\}). \end{aligned} \tag{2.38}$$

Here we used

$$\sum_{j,k=1}^n \int_{\mathbb{R}^n} |x|^{-2} x_j x_k \overline{f(x)} f_{x_j,x_k}(x)\, d^n x = -\int_{\mathbb{R}^n} |x|^{-2} |[x\cdot(\nabla f)(x)]|^2\, d^n x \\ - (n-1)\int_{\mathbb{R}^n} |x|^{-2} \overline{f(x)} [x\cdot(\nabla f)(x)]\, d^n x, \quad f\in C_0^\infty(\mathbb{R}^n\backslash\{0\}). \tag{2.39}$$

Thus,

$$\int_{\mathbb{R}^n} \big|\big[|x|^{-1} x\cdot\nabla f\big](x)\big|^2\, d^n x \geq \alpha[(n-2)-\alpha]\int_{\mathbb{R}^n} |x|^{-2}|f(x)|^2\, d^n x. \tag{2.40}$$

Maximizing $\alpha[(n-2)-\alpha]$ with respect to α yields the improved Hardy inequality,

$$\int_{\mathbb{R}^n} \big|\big[|x|^{-1} x\cdot\nabla f\big](x)\big|^2\, d^n x \geq [(n-2)/2]^2\int_{\mathbb{R}^n} |x|^{-2}|f(x)|^2\, d^n x. \tag{2.41}$$

(By Cauchy's inequality, (2.41) implies the classical Hardy inequality (2.32).) Again, it is known that the constant in (2.41) is optimal (cf., e.g., [6, Theorem 1.2.5]). ⋄

Remark 2.10. The case of Rellich (and Hardy) inequalities in the half-line case is completely analogous (and much more straightforward): Consider the differential expressions

$$T = -\frac{d^2}{dx^2} + \frac{\alpha}{x}\frac{d}{dx} + \frac{\beta}{x^2}, \quad T^+ = -\frac{d^2}{dx^2} - \frac{\alpha}{x}\frac{d}{dx} + \frac{\alpha+\beta}{x^2}, \tag{2.42}$$

with $\alpha,\beta\in\mathbb{R}$, which are formal adjoints to each other. One verifies,

$$T^+T = \frac{d^4}{dx^4} + \frac{\alpha-\alpha^2-2\beta}{x^2}\frac{d^2}{dx^2} + \frac{2\alpha^2-2\alpha+4\beta}{x^3}\frac{d}{dx} + \frac{3\alpha\beta+\beta^2-6\beta}{x^4}, \tag{2.43}$$

and hence upon some integrations by parts,

$$0 \leq \int_0^\infty (Tf)(x)^2\, dx = \int_0^\infty f(x)(T^+Tf)(x)\, dx \\ = \int_0^\infty [f''(x)]^2\, dx - \big(\alpha-\alpha^2-2\beta\big)\int_0^\infty \frac{[f'(x)]^2}{x^2}\, dx \\ + \beta(3\alpha+\beta-6)\int_0^\infty \frac{f(x)^2}{x^4}\, dx, \quad f\in C_0^\infty((0,\infty)), \tag{2.44}$$

choosing f real-valued (for simplicity and w.l.o.g.). Thus, one obtains,

$$\int_0^\infty |f''(x)|^2\, dx \geq \big(\alpha-\alpha^2-2\beta\big)\int_0^\infty \frac{|f'(x)|^2}{x^2}\, dx \\ + \beta(6-\beta-3\alpha)\int_0^\infty \frac{|f(x)|^2}{x^4}\, dx, \\ f\in C_0^\infty((0,\infty)),\ \alpha,\beta\in\mathbb{R}. \tag{2.45}$$

Choosing $\beta = \big(\alpha-\alpha^2\big)/2$ yields the Rellich-type inequality

$$\int_0^\infty |f''(x)|^2\, dx \geq \big[3\alpha - (19/4)\alpha^2 + 2\alpha^3 - (1/4)\alpha^4\big]\int_0^\infty \frac{|f(x)|^2}{x^4}\, dx, \\ f\in C_0^\infty((0,\infty)). \tag{2.46}$$

Introducing $F(\alpha) = 3\alpha - (19/4)\alpha^2 + 2\alpha^3 - (1/4)\alpha^4 = -(1/4)(\alpha-4)(\alpha-3)(\alpha-1)\alpha$, $\alpha\in\mathbb{R}$, one verifies that $F(2+\gamma) = F(2-\gamma)$, $\gamma\in\mathbb{R}$, and factors its derivative as

$$F'(\alpha) = 3 - (19/2)\alpha + 6\alpha^2 - \alpha^3 \\ = -(\alpha-2)\big(\alpha-2+(5/2)^{1/2}\big)\big(\alpha-2-(5/2)^{1/2}\big). \tag{2.47}$$

One notes that $\alpha_1 = 2$ yields a local minimum with $F(2) = -1$, $\alpha_2 = 2 - (5/2)^{1/2}$ and $\alpha_3 = 2 + (5/2)^{1/2}$ both yield local maxima of equal value, that is, $F(\alpha_2) = F(\alpha_3) = \frac{9}{16}$. Thus, one obtains Rellich's inequality for the half-line in the form

$$\int_0^\infty |f''(x)|^2 \, dx \geq \frac{9}{16} \int_0^\infty \frac{|f(x)|^2}{x^4} \, dx, \quad f \in C_0^\infty((0,\infty)). \tag{2.48}$$

We refer to Birman [13, p. 46] (see also Glazman [34, p. 83–84]), who presents a sequence of higher-order Hardy-type inequalities on $(0,\infty)$ whose second member coincides with (2.48). For a variant of (2.48) on the interval $(0,1)$ we refer to [16, p. 114]; the case of higher-order Hardy-type inequalities for general interval is also considered in [52]. We will reconsider this sequence of higher-order Hardy-type inequalities in [30].

In addition, choosing $\beta = 0$ or $\beta = 6 - 3\alpha$ and subsequently maximizing with respect to α yields in either case

$$\int_0^\infty |f''(x)|^2 \, dx \geq \frac{1}{4} \int_0^\infty \frac{|f'(x)|^2}{x^2} \, dx, \quad f \in C_0^\infty((0,\infty)), \tag{2.49}$$

however, this is just Hardy's inequality [36], [37], with f replaced by f'. $\diamond$

Remark 2.11. While we basically focused on inequalities in $L^2(\mathbb{R}^n)$ (see, however, Remark 2.6), much of the recent work on Rellich and higher-order Hardy inequalities aims at $L^p(\Omega)$ for open sets $\Omega \subset \mathbb{R}^n$ (frequently, Ω is bounded with $0 \in \Omega$), $p \in [1,\infty)$, appropriate remainder terms (the latter often associated with logarithmic refinements or with boundary terms), higher-order Hardy–Rellich inequalities, and the inclusion of magnetic fields. The enormous number of references on this subject, especially, in the context of Hardy-type inequalities, makes it impossible to achieve any reasonable level of completeness in such a short note as the underlying one. Hence we felt we had to restrict ourselves basically to Rellich and higher-order Hardy inequality references only and thus we refer, for instance, to [1], [2], [4], [5], [6, Ch. 6], [7], [8], [9], [11], [12], [17], [18], [19], [20], [22], [23], [24], [25], [26], [27], [33], [40], [45], [46], [48], [49], [52], [53], [55], [58], [59], [60], and the extensive literature cited therein. For the case of Hardy-type inequalities we only refer to the standard monographs such as, [6], [43], [44], and [51].

In this context we emphasize once again that the factorization method is entirely independent of the choice of domain Ω. Indeed, factorizations in the context of Hardy's inequality in balls with optimal constants and logarithmic correction terms were already studied in [28], [31], based on prior work in [38], [41], and [42], although this appears to have gone unnoticed in the recent literature on this subject. For instance, one can introduce iterated logarithms of the form for $\gamma > 0$, $x \in \mathbb{R}^n$, $|x| < \gamma$,

$$\begin{aligned} &(-\ln(|x|/\gamma))_0 = 1, \\ &(-\ln(|x|/\gamma))_1 = (-\ln(|x|/\gamma)), \\ &(-\ln(|x|/\gamma))_{k+1} = \ln((-\ln(|x|/\gamma))_k), \quad k \in \mathbb{N}, \end{aligned} \tag{2.50}$$

and replace ∇ by $T_{\alpha_m,y}$, where

$$T_{\alpha_m,y} = \nabla + 2^{-1}|x-y|^{-2}\bigg\{(n-2) + \sum_{j=1}^{m}\prod_{k=1}^{j}[(-\ln(|x-y|/\gamma))_k]^{-1} - \alpha_m \prod_{k=1}^{m}[(-\ln(|x-y|/\gamma))_k]^{-1}\bigg\}(x-y), \quad 0<|x|<r,\ r<\gamma,\ \alpha_m \geq 0,\ m\in\mathbb{N}, \tag{2.51}$$

$$T_{\alpha_0,y} = \nabla + 2^{-1}(n-2-\alpha_0)|x-y|^{-2}(x-y), \quad 0<|x|<r,\ \alpha_0\geq 0,\ m=0. \tag{2.52}$$

Then with $T^+_{\alpha_m,y}$ the formal adjoint of $T_{\alpha_m,y}$, one obtains for $f\in C_0^\infty(B_n(y;r)\backslash\{y\})$

$$(T^+_{\alpha_m,y}T_{\alpha_m,y}f)(x) = (-\Delta f)(x) - 4^{-1}|x-y|^{-2}\bigg\{(n-2)^2 + \sum_{j=1}^{m}\sum_{k=1}^{j}[(-\ln(|x-y|/\gamma))_k]^{-2} f(x) - \alpha_m^2\prod_{k=1}^{m}[(-\ln(|x-y|/\gamma))_k]^{-2}\bigg\}f(x), \quad m\in\mathbb{N}, \tag{2.53}$$

$$(T^+_{\alpha_0,y}T_{\alpha_0,y}f)(x) = (-\Delta f)(x) - 4^{-1}\big[(n-2)^2-\alpha_0^2\big]|x-y|^{-2}f(x), \quad m=0. \tag{2.54}$$

(Here $B_n(x_0;r_0)$ denotes the open ball in $\mathbb{R}^n$ with center $x_0\in\mathbb{R}^n$ and radius $r_0>0$.) In particular, letting $r_0\downarrow 0$ and $r_1\uparrow r$ in [28, Lemma 1] implies

$$0\leq \int_{B(y;r)}|(T_{\alpha_m,y}f)(x)|^2 = \int_{B(y;r)}\bigg\{|(\nabla f)(x)|^2 - 4^{-1}|x-y|^{-2}\bigg[(n-2)^2 + \sum_{j=1}^{m}\sum_{k=1}^{j}[(-\ln(|x-y|/\gamma))_k]^{-2} - \alpha_m^2\prod_{k=1}^{m}[(-\ln(|x-y|/\gamma))_k]^{-2}\bigg]|f(x)|^2\bigg\}\,d^nx, \quad 0<r<\gamma,\ f\in C_0^\infty(B(y;r)\backslash\{y\}),\ m\in\mathbb{N}\cup\{0\}, \tag{2.55}$$

and hence (with $\alpha_m=0$),

$$\int_{B(y;r)}|(\nabla f)(x)|^2\,d^nx \geq 4^{-1}\int_{B(y;r)}|x-y|^{-2}\bigg\{(n-2)^2 + \sum_{j=1}^{m}\sum_{k=1}^{j}[(-\ln(|x-y|/\gamma))_k]^{-2}\bigg\}|f(x)|^2\,d^nx, \quad 0<r<\gamma,\ f\in C_0^\infty(B(y;r)\backslash\{y\}),\ m\in\mathbb{N}\cup\{0\}. \tag{2.56}$$

(Following standard practice, a product, resp., sum over an empty index set is defined to equal 1, resp., 0.) In analogy to Remark 2.6, inequality (2.56) extends to arbitrary open bounded sets $\Omega\subset\mathbb{R}^n$ as long as γ is chosen sufficiently large (e.g., larger than the diameter of Ω). The constants in (2.55) are best possible as it is well-known that the operators (2.53), (2.54) are nonnegative if and only if $\alpha_m^2\geq 0$, $m\in\mathbb{N}\cup\{0\}$ (cf., e.g., [28, p. 99] or [32, Theorem 2.2]). (They are unbounded from below for $\alpha_m^2<0$, $m\in\mathbb{N}\cup\{0\}$, permitting temporarily a continuation to negative values of α^2 on the right-hand sides of (2.53) and (2.54).)

For the special half-line case we also refer to [32].

Higher-order logarithmic refinements of the multi-dimensional Hardy–Rellich-type inequality appeared in [1, Theorem 2.1], and a sequence of such multi-dimensional Hardy–Rellich-type inequalities, with additional generalizations, appeared in [58, Theorems 1.8–1.10]. ⋄

We conclude by mentioning that factorization also works for other singular interactions, for instance, for point dipole interactions, where $|x|^{-2}$ is replaced by $|x|^{-3}(d\cdot x)$, with $d \in \mathbb{R}^n$ a constant vector. Moreover, it applies to higher-order Hardy-type inequalities where $-\Delta$ is replaced by $(-\Delta)^\ell$, $\ell \in \mathbb{N}$. We defer all this to future investigations [29].

Acknowledgments. We are indebted to Mark Ashbaugh and Roger Lewis for very valuable hints to the literature on Hardy–Rellich-type inequalities and to Richard Wellmann for helpful discussions.

References

[1] Adimurthi, M. Grossi, and S. Santra, *Optimal Hardy–Rellich inequalities, maximum principle and related eigenvalue problem*, J. Funct. Anal. **240**, 36—83 (2006).
[2] W. Allegretto, *Nonoscillation theory of elliptic equations of order 2n*, Pacific J. Math. **64**, 1–16 (1976).
[3] W. Arendt, G. R. Goldstein, J. A. Goldstein, *Outgrowths of Hardy's inequality*, in *Recent Advances in Differential Equations and Mathematical Physics*, N. Chernov, Y. Karpeshina, I. W. Knowles, R. T. Lewis, and R. Weikard (eds.), Contemp. Math. **412**, 51–68, 2006.
[4] F. G. Avkhadiev, *Rellich type inequalities in domains of the Euclidean space*, Russ. Math. **60**, 60–63 (2016).
[5] F. G. Avkhadiev, *Hardy–Rellich inequalities in domains of the Euclidean space*, J. Math. Anal. Appl. **442**, 469–484 (2016).
[6] A. A. Balinsky, W. D. Evans, and R. T. Lewis, *The Analysis and Geometry of Hardy's Inequality*, Universitext, Springer, 2015.
[7] G. Barbatis, *Improved Rellich inequalities for the polyharmonic operator*, Indiana Univ. Math. J. **55**, 1401–1422 (2006).
[8] G. Barbatis, *Best constants for higher-order Rellich inequalities*, Math. Z. **255**, 877–896 (2007).
[9] G. Barbatis and A. Tertikas, *On a class of Rellich inequalities*, J. Comp. Appl. Math. **194**, 156–172 (2006).
[10] D. M. Bennett, *An extension of Rellich's inequality*, Proc. Amer. Math. Soc. **106**, 987–993 (1989).
[11] E. Berchio, *On some higher order Hardy–Rellich type inequalities with boundary terms*, Nonlinear Anal. **70**, 2834–2841 (2009).
[12] E. Berchio, D. Cassani, and F. Gazzola, *Hardy–Rellich inequalities with boundary remainder terms and applications*, Manuscripta Math. **131**, 427–458 (2010).
[13] M. S. Birman, *The spectrum of singular boundary problems*, Amer. Math. Soc. Transl., Ser. 2, **53**, 23–80 (1966).
[14] D. G. Costa, *On Hardy–Rellich type inequalities in* $\mathbb{R}^N$, Appl. Math. Lett. **22**, 902–905 (2009).
[15] E. B. Davies, *Heat Kernels and Spectral Theory*, Cambridge Tracts in Math., Vol. 92, Cambridge Univ. Press, Cambridge, 1989.
[16] E. B. Davies, *Spectral Theory and Differential Operators*, Cambridge University Press, Cambridge, 1995.
[17] E. B. Davies and A. M. Hinz, *Explicit constants for Rellich inequalities in* $L_p(\Omega)$, Math. Z. **227**, 511–523 (1998).
[18] A. Detalla, T. Horiuchi, and H. Ando, *Missing terms in Hardy–Sobolev inequalities*, Proc. Japan Acad. **80A**, 160–165 (2004).
[19] A. Detalla, T. Horiuchi, and H. Ando, *Missing terms in Hardy–Sobolev inequalities and its applications*, Far East J. Math. Sci. **14**, 333–359 (2004).
[20] A. Detalla, T. Horiuchi, and H. Ando, *Sharp remainder terms of the Rellich inequality and its applications*, Bull. Malaysian Math. Sci. Soc. **35**, 519–528 (2012).
[21] D. E. Edmunds and W. D. Evans, *Spectral Theory and Differential Operators*, Clarendon Press, Oxford, 1989.
[22] D. E. Edmunds and W. D. Evans, *The Rellich inequality*, Rev. Mat. Complut. **29**, 511–530 (2016).
[23] W. D. Evans, *Recent results on Hardy and Rellich inequalities*, in *Further Progress in Analysis*, H. G. W. Begehr, A. O. Gelebi, R. P. Gilbert, and H. T. Kaptanoğlu (eds.), World Scientifc, New Jersey, 2009, pp. 33–44.
[24] W. D. Evans and R. T. Lewis, *On the Rellich inequality with magnetic potentials*, Math. Z. **251**, 267–284 (2005).
[25] W. D. Evans and R. T. Lewis, *Hardy and Rellich inequalities with remainders*, J. Math. Inequal. **1**, 473–490 (2007).

[26] V. A. Galaktionov, *On extensions of higher-order Hardy's inequalities*, Diff. Integral Eq. **19**, 327–344 (2006).
[27] F. Gazzola, H.-C. Grunau, and E. Mitidieri, *Hardy inequalities with optimal constants and remainder terms*, Trans. Amer. Math. Soc. **356**, 2149–2168 (2003).
[28] F. Gesztesy, *On non-degenerate ground states for Schrödinger operators*, Rep. Math. Phys. **20**, 93–109 (1984).
[29] F. Gesztesy, I. Michael, and L. Littlejohn, in preparation.
[30] F. Gesztesy, I. Michael, L. Littlejohn, and R. Wellman, *A note on Birman's sequence of Hardy–Rellich-type inequalities*, in preparation.
[31] F. Gesztesy and L. Pittner, *A generalization of the virial theorem for strongly singular potentials*, Rep. Math. Phys. **18**, 149–162 (1980).
[32] F. Gesztesy and M. Ünal, *Perturbative oscillation criteria and Hardy-type inequalities*, Math. Nachr. **189**, 121–144 (1998).
[33] N. Ghoussoub and A. Moradifam, *Bessel pairs and optimal Hardy and Hardy–Rellich inequalities*, Math. Ann. **349**, 1–57 (2011).
[34] I. M. Glazman, *Direct Methods of Qualitative Spectral Analysis of Singular Differential Operators*, Israel Program for Scientific Translations, Jerusalem, 1965.
[35] J. A. Goldstein and R. Svirsky, *On a domain characterization of Schrödinger operators with gradient magnetic vector potentials and singular potentials*, Proc. Amer. Math. Soc. **105**, 317–323 (1989).
[36] H. Hardy, *Notes on some points in the integral calculus*, Messenger Math. **48**, 107–112 (1919).
[37] G. H. Hardy, *Note on a theorem of Hilbert*, Math. Z. **6**, 314–317 (1920).
[38] H. Kalf, *On the characterization of the Friedrichs extension of ordinary or elliptic differential operators with a strongly singular potential*, J. Funct. Anal. **10**, 230–250 (1972).
[39] H. Kalf, *Gauss' theorem and the self-adjointness of Schrödinger operators*, Arkiv Mat. **18**, 19–47 (1980).
[40] H. Kalf, *A note on the domain characterization of certain Schrödinger operators with strongly singular potentials*, Proc. Roy. Soc. Edinburgh **97A**, 125–130 (1984).
[41] H. Kalf, U.-W. Schmincke, J. Walter, and R. Wüst, *On the spectral theory of Schrödinger and Dirac operators with strongly singular potentials*, in *Spectral Theory and Differential Equations*, W. N. Everitt (ed.), Lecture Notes in Math., Vol. 448, Springer, Berlin, 1975, pp. 182–226.
[42] H. Kalf and J. Walter, *Strongly singular potentials and essential self-adjointness of singular elliptic operators in $C_0^\infty(\mathbb{R}^n\backslash\{0\})$*, J. Funct. Anal. **10**, 114–130 (1972).
[43] A. Kufner, L. Maligranda, and L.-E. Persson, *The Hardy Inequality. About its History and Some Related Results*, Vydavatelský Servis, Pilsen, 2007.
[44] A. Kufner and L.-E. Persson, *Weighted Inequalities of Hardy Type*, World Scientific, Singapore, 2003.
[45] G. Metafune, M. Sobajima, and C. Spina, *Weighted Calderón–Zygmund and Rellich inequalities in L^p*, Math. Ann. **361**, 313–366 (2015).
[46] G. Metafune, M. Sobajima, and C. Spina, *Rellich and Calderón–Zygmund inequalities for an operator with discontinuous coefficients*, Ann. Mat. Pura Appl. **195**, 1305–1331 (2016).
[47] E. Mitidieri, *A simple approach to Hardy inequalities*, Math. Notes **67**, 479–486 (2000).
[48] A. Moradifam, *Optimal weighted Hardy–Rellich inequalities on $H^2 \cap H_0^1$*, J. Lond. Math. Soc. (2) **85**, 22–40 (2012).
[49] R. Musina, *Optimal Rellich–Sobolev constants and their extremals*, Diff. Integral Eq. **27**, 579–600 (2014).
[50] N. Okazawa, L^p-theory of Schrödinger operators with strongly singular potentials, Japan J. Math. **22**, 199–239 (1996).
[51] B. Opic and A. Kufner, *Hardy-Type Inequalities*, Pitman Research Notes in Mathematics Series, Vol. 219. Longman Scientific & Technical, Harlow, 1990.
[52] M. P. Owen, *The Hardy–Rellich inequality for polyharmonic operators*, Proc. Roy. Soc. Edinburgh **A 129**, 825–839 (1999).
[53] B. G. Pachpatte, *On some extensions of Rellich's inequality*, Tamkang J. Math. **22**, 259–265 (1991).
[54] F. Rellich, *Perturbation Theory of Eigenvalue Problems*, Gordon and Breach, New York, 1969.
[55] M. Ruzhansky and D. Suragan, *Hardy and Rellich inequalities, and sharp remainders on homogeneous groups*, arXiv: 1603.06239.
[56] U-W. Schmincke, *Essential self-adjointness of a Schrödinger operator with strongly singular potential*, Math. Z. **124**, 47–50 (1972).
[57] B. Simon, *Hardy and Rellich inequalities in non-integral dimension*, J. Operator Th. **9**, 143–146 (1983). Addendum, J. Opertor Th. **12**, 197 (1984).
[58] A. Tertikas and N. B. Zographopoulos, *Best constants in the Hardy–Rellich inequalities and related improvements*, Adv. Math. **209**, 407–459 (2007).

[59] C. Xia, *Hardy and Rellich type inequalities on complete manifolds*, J. Math. Anal. **409**, 84–90 (2014).
[60] Y.-X. Xiao and Q.-H. Yang, *An improved Hardy–Rellich inequality with optimal constant*, J. Inequalities Appls. **2009**, Art. ID 610530, 10 pp.
[61] D. Yafaev, *Sharp constants in the Hardy–Rellich inequalities*, J. Funct. Anal. **168**, 121–144 (1999).

Department of Mathematics, Baylor University, One Bear Place #97328, Waco, TX 76798-7328, USA
E-mail address: Fritz_Gesztesy@baylor.edu
URL: http://www.baylor.edu/math/index.php?id=935340

Department of Mathematics, Baylor University, One Bear Place #97328, Waco, TX 76798-7328, USA
E-mail address: Lance_Littlejohn@baylor.edu
URL: http://www.baylor.edu/math/index.php?id=53980